

References using FD Neutral Red Solution™ (PS106-01 or PS106-02):

1. Calabrese E, Du F, Garman RH, Johnson GA, Riccio C, Tong LC, Long JB. (2014) Diffusion tensor imaging reveals white matter injury in a rat model of repetitive blast-induced traumatic brain injury. **J Neurotrauma**. 31(10):938-50.
2. Parr-Brownlie LC, Poloskey SL, Bergstrom DA, and Walters JR. (2009) Parafascicular thalamic nucleus activity in a rat model of Parkinson's disease. **Exp Neurol**. 217(2):269-81.
3. Parr-Brownlie LC, Poloskey SL, Flanagan KK, Eisenhofer G, Bergstrom DA, and Walters JR. (2007) Dopamine lesion-induced changes in subthalamic nucleus activity are not associated with alterations in firing rate or pattern in layer V neurons of the anterior cingulate cortex in anesthetized rats. **Eur. J Neurosci**. 26(7):1925-39.
4. Behl B, Klos M, Serr M, Ebert U, Janson B, Drescher K, Gross G, and Schoemaker H. (2006) An ELISA-based method for the quantification of incorporated BrdU as a measure of cell proliferation in vivo. **J Neurosci Meth**. 158(1):37-49.
5. Rakhade SN, Yao B, Ahmed S, Asano E, Beaumont TL, Shah AK, Draghici S, Krauss R, Chugani HT, Sood S, and Loeb JA. (2005) A common pattern of persistent gene activation in human neocortical epileptic foci. **Ann Neurol**. 58(5):736-47.